

10-217

**PROCESSED BY**

NOV 4 - 1998

**FBI**

In re Application of

*Queen et al*

Application Number

310,252

Filed

2/13/89

Group Art Unit

Examiner

Paper No.

*#22*

Assistant Commissioner for Patents  
Washington, DC 20231

I hereby request access under 37 CFR 1.14(a)(3)(iv) to the application file record of the above-identified **ABANDONED** application, which is: (CHECK ONE)

☒ (A) referred to in United States Patent Number 5693761, column \_\_\_\_\_

\_\_\_\_ (B) referred to in an application that is open to public inspection as set forth in 37 CFR 1.11, i.e., Application No. \_\_\_\_\_, filed \_\_\_\_\_, on page \_\_\_\_\_ of paper number \_\_\_\_\_

\_\_\_\_ (C) an application that claims the benefit of the filing date of an application that is open to public inspection, i.e., Application No. \_\_\_\_\_, filed \_\_\_\_\_, or

\_\_\_\_ (D) an application in which the applicant has filed an authorization to lay open the complete application to the public.

Please direct any correspondence concerning this request to the following address:

\_\_\_\_\_  
\_\_\_\_\_  
\_\_\_\_\_

*Nancy Perry*  
\_\_\_\_\_  
Signature

*Nancy Perry*  
\_\_\_\_\_  
Typed or printed name

Date

FOR PTO USE ONLY

Approved by: *[Signature]*

(initials)

Unit: *F.I.U.*



US005693761A

10-217

**United States Patent** [19]

Queen et al.

[11] Patent Number: 5,693,761

[45] Date of Patent: Dec. 2, 1997

[54] POLYNUCLEOTIDES ENCODING  
IMPROVED HUMANIZED  
IMMUNOGLOBULINS[75] Inventors: Cary L. Queen, Los Altos; William P.  
Schneider, Mountain View; Harold E.  
Selick, Belmont, all of Calif.[73] Assignee: Protein Design Labs, Inc., Mountain  
View, Calif.

[21] Appl. No.: 474,040

[22] Filed: Jun. 7, 1995

**Related U.S. Application Data**[62] Division of Ser. No. 634,278, Dec. 19, 1990, Pat. No.  
5,530,101, which is a continuation of Ser. No. 590,274, Sep.  
28, 1990, abandoned, and a continuation of Ser. No. 310,  
252, Feb. 13, 1989, abandoned, which is a continuation of  
Ser. No. 290,975, Dec. 28, 1988, abandoned.[51] Int. Cl.<sup>6</sup> ..... C07H 21/04[52] U.S. Cl. .... 536/23.53; 530/387.3;  
435/320.1; 435/252.3[58] Field of Search ..... 536/23.53; 530/387.3;  
435/320.1, 252.3[56] **References Cited****U.S. PATENT DOCUMENTS**

4,578,335	3/1986	Urdal et al.	530/351
4,816,397	3/1989	Boss et al.	435/68
4,816,565	3/1989	Honjo et al.	435/69.1
4,816,567	3/1989	Cabilly et al.	530/387
4,845,198	7/1989	Urdal et al.	530/387
4,867,973	9/1989	Goers et al.	424/85.91
5,198,359	3/1993	Taniguchi et al.	435/252.3
5,225,539	7/1993	Winter	530/387.3
5,476,786	12/1995	Huston et al.	435/85.8

**FOREIGN PATENT DOCUMENTS**

0 120 694	10/1984	European Pat. Off.
0171496	2/1986	European Pat. Off.
0173494	3/1986	European Pat. Off.
0184187	6/1986	European Pat. Off.
0256654	7/1987	European Pat. Off.
0239400	9/1987	European Pat. Off.
0266663	6/1988	European Pat. Off.
0 323 806	7/1989	European Pat. Off.
0 328 404	8/1989	European Pat. Off.
0 365 209	4/1990	European Pat. Off.
0 365 997	5/1990	European Pat. Off.
0 125 023	6/1991	European Pat. Off.
0460167	12/1991	European Pat. Off.
2188941	10/1987	United Kingdom
8928874	12/1989	United Kingdom
WO 86/05513	9/1986	WIPO
WO 87/02671	5/1987	WIPO
WO 88/09344	12/1988	WIPO
WO 89/01783	3/1989	WIPO
91/09967	7/1991	WIPO

**OTHER PUBLICATIONS**Better et al., "Escherichia coli Secretion of an Active Chi-  
meric Antibody Fragment," *Science*, 240:1041-1043  
(1988).Bird et al., "Single-Chain Antigen-Binding Proteins," *Sci-  
ence*, 242:423-426 (1988).Boulianne et al., "Production of functional chimeric mouse/  
human antibody," *Nature*, 312:643-646 (1984).Carter et al., "Humanization of an anti-p185<sup>HER2</sup> antibody  
for human cancer therapy," *Proc. Natl. Acad. Sci.*,  
89:4285-4289 (1992).Chothia, C. and A.M. Lesk, "Canonical Structures for the  
Hypervariable Regions of Immunoglobulins," *J. Mol. Biol.*,  
196:901-917 (1987).Co et al., "Humanized antibodies for antiviral therapy,"  
*Proc. Natl. Acad. Sci.*, 88:2869-2873 (1991).Co et al., "Chimeric and humanized antibodies with speci-  
ficity for the CD33 antigen," *J. Immunol.*, 148:1149-1154  
(1992).Daugherty et al., "Polymerase chain reaction facilitates the  
cloning, CDR-grafting, and rapid expansion of a murine  
monoclonal antibody directed against the CD18 component  
of leukocyte integrins," *Nuc. Acids. Res.*, 19:2471-2476  
(1991).Ellison et al., "The nucleotide sequence of a human immu-  
noglobulin c(gamma)<sub>1</sub> gene," *Nucleic Acids Res.*, 10:4071-  
(1982).Farrar, J., "The biochemistry, biology, and the role of  
interleukin-2 in the induction of cytotoxic T cell and anti-  
body-forming B cell receptors," *Immunol. Rev.*, 63:129-166  
(1982).Foote et al., "Antibody framework residues affecting the  
conformation of hypervariable loops," *J. Mol. Biol.*,  
224:487-499 (1992).Gorman et al., "Reshaping a therapeutic CD4 antibody,"  
*Proc. Natl. Acad. Sci.*, 88:4181-4185 (1991).

(List continued on next page.)

Primary Examiner—Lila Feisee

Assistant Examiner—Julie E. Reeves

Attorney, Agent, or Firm—Townsend and Townsend and  
Crew LLP[57] **ABSTRACT**

Novel methods for producing, and compositions of, human-  
ized immunoglobulins having one or more complementarity  
determining regions (CDR's) and possible additional amino  
acids from a donor immunoglobulin and a framework region  
from an accepting human immunoglobulin are provided.  
Each humanized immunoglobulin chain will usually  
comprise, in addition to the CDR's, amino acids from the  
donor immunoglobulin framework that are, e.g., capable of  
interacting with the CDR's to effect binding affinity, such as  
one or more amino acids which are immediately adjacent to  
a CDR in the donor immunoglobulin or those within about  
about 3 Å as predicted by molecular modeling. The heavy  
and light chains may each be designed by using any one or  
all of various position criteria. When combined into an intact  
antibody, the humanized immunoglobulins of the present  
invention will be substantially non-immunogenic in humans  
and retain substantially the same affinity as the donor  
immunoglobulin to the antigen, such as a protein or other  
compound containing an epitope.

37 Claims, 55 Drawing Sheets

It is time to start looking more deeply for common attributes of all the immortalizing genes. Do others, apart from large T and p53, interact with each other and proceed through a common pathway? That some sophistication in our outlook is needed is highlighted by the recent characterization of an Abelson virus-transformed cell line, L12, that does not express p53 because its gene has been inactivated by insertion of a retrovirus-like sequence into its first intron<sup>12</sup>. L12 cells grow continuously in culture but, unlike other Abelson virus-transformed cells, will not give rise to lethal tumours in syngenic animals. Instead the tumours regress spontaneously despite reaching considerable size. When a functional p53 gene is reintroduced into L12 cells, those cells which then express the gene are able to give rise to progressive and lethal tumours. So, in this system, the p53 gene product seems to act as a tumour progression factor, whose effect is only really detectable in an *in vivo* assay. This should delight rather than mystify, because the

existence of a battery of oncogenes of subtly different effects, gives us a better chance to simplify and categorize the apparent plethora of human neoplastic diseases. □

1. Parada, L.F., Land, H., Weinberg, R.A., Wolf, D. & Rotter, V. *Nature* 312, 649 (1984).
2. Eliyahu, D., Raz, A., Gruss, P., Givol, D. & Oren, M. *Nature* 312, 646 (1984).
3. Jenkins, J.R., Rudge, K. & Currie, G.A. *Nature* 312, 651 (1984).
4. Lane, D.P. & Crawford, L.V. *Nature* 278, 261 (1979).
5. Sarnow, P., Ho, Y.S., Williams, J. & Levine, A.J. *Cell* 28, 387 (1982).
6. Crawford, L.V., Pim, D.C., Gurney, E.G., Goodfellow, P. & Taylor-Papadimitrou, J. *Proc. natn. Acad. Sci. U.S.A.* 78, 41 (1981).
7. DeLeo, A.B., et al. *Proc. natn. Acad. Sci. U.S.A.* 76, 2420 (1979).
8. Mercer, W.E., Nelson, D., DeLeo, A.B., Old, L.J. & Baserga, R. *Proc. natn. Acad. Sci. U.S.A.* 79, 6309 (1982).
9. Ruley, H.E. *Nature* 304, 602 (1983).
10. Land, H., Parada, L.F. & Weinberg, R.A. *Nature* 304, 596 (1983).
11. Colby, W.W. & Shenk, T. *Proc. natn. Acad. Sci. U.S.A.* 79, 5189 (1982).
12. Wolf, D., Harris, N. & Rotter, V. *Cell* 38, 119 (1984).

D.P. Lane is in the Cancer Research Campaign Eukaryotic Genetics Research Group, Imperial College, London SW7 2AZ, UK.

## Immunology

# Uses of chimaeric antibodies

from Alan Munro

ANTIBODIES were discovered through their ability to neutralize bacterial toxins, and for almost one hundred years antitoxin sera from experimental animals have been used to save many lives. For a brief period, it seemed that serum therapy would solve numerous medical problems. But it was soon found that repeated injections of a supposedly innocuous 'foreign' antibody can lead to harmful hypersensitivity reactions. With the availability of monoclonal antibodies, many new applications of serum therapy (for example, see last week's *Nature*; Cobbold *et al.* 312, 548) would become well worth exploring if only hypersensitivity reactions could be avoided.


There are three ways to approach this problem. The first is to use monoclonal antibodies from the standard source — hybridomas of rat or mouse cells — but to induce specific unresponsiveness to rodent immunoglobulin before or during therapy. The second approach is to use human monoclonal antibodies, produced either from human hybridomas or from Epstein-Barr virus-immortalized human B lymphocyte cell lines; despite considerable efforts, the success of this approach has been limited and there is no generalized way of obtaining human monoclonal antibodies of the right specificity. Finally, by using the techniques of genetic engineering, it may be possible to obtain antibodies in which the antigen-binding site is defined by sequences from a rodent monoclonal antibody of the right specificity whereas the rest of the molecule is as 'human' in structure as possible.

Papers by G.L. Boulianne, N. Hozumi and M.J. Shulman on page 643 of this issue of *Nature* and by S.L. Morrison *et al.* in *Proc. natn. Acad. Sci. U.S.A.* 81, 6851; 1984 are the first to describe functional chimaeric antibody molecules containing light chains and heavy chains, in which the whole of the variable regions of the chains are 'mouse' and the constant regions are 'human'. Possibly the human immune system will not recognize such chimaeric molecules as foreign, but there are good reasons to think that it will. In many cases it is possible to mount an immune response against antibody molecules themselves, even when the only antigenic determinants lie in the variable regions (take, for example, the heavy-chain variable region allotypes in rabbits). At least it may be easier to induce unresponsiveness to rodent variable regions when they are part of chimaeric antibodies. In that respect, it will be interesting to compare the rates of clearance of chimaeric molecules with those of the normal antibodies molecules of the host. Even if it proves possible to construct chimaeric antibodies with rodent hypervariable regions embedded in a human variable region framework, it is likely that they will be recognized by the human immune system — but the hope is that such chimaeras will be sufficiently 'self-like' to become integrated into the network of interacting idiotypes and anti-idiotypes without sensitizing the host and so avoiding subsequent hypersensitivity reactions.

The construction of chimaeric antibodies has been made possible through the

development, by a number of independent groups, of vector systems that allow the expression of antibody light-chain or heavy-chain genes in cells that have been transfected with them. Many laboratories have since initiated experiments in which cells are transfected with modified antibody genes. Earlier this year J. Sharon *et al.* (*Nature* 309, 364; 1984) reported the expression of a novel fragment of antibody consisting of a normal light chain ( $V_L C_\kappa$ ) associated with a modified antibody chain made up of a heavy-chain variable region linked to a light-chain constant region ( $V_H C_\lambda$ ). This fragment binds antigen and is secreted from the transfected cells (showing that the Fc portion of antibody is not necessary for secretion).

The report by M.S. Neuberger, G.T. Williams and R.O. Fox on page 604 of this issue not only confirms that result but goes much further. It shows that the Fc portion of the antibody heavy chain can be replaced by a completely unrelated protein — the enzyme staphylococcal nuclease for example — to produce a bifunctional



Schematic diagram of a mouse antibody showing two identical light chains (L) and two identical heavy chains (H). Each chain consists of variable (V) and constant (C) domains. The antibody class is determined by the heavy-chain constant region which imparts distinctive features on the tail (Fc) region. Neuberger *et al.* have substituted staphylococcal nuclease for the Fc region of both heavy chains of immunoglobulin M; Boulianne *et al.* have replaced the C domain of each heavy chain of the immunoglobulin M monomer with a human C domain.

chimaeric molecule — in this case, exhibiting both specific antigen-binding and nuclease activity. There are many potential applications of this technology in the purification of cloned gene products, in enzyme-linked immunoassays and in the targeting of chemotherapeutic drugs on to tumour cells. □

Alan Munro is in the Department of Pathology, University of Cambridge, Tennis Court Road, Cambridge CB2 1OP, UK.